

Exercise Problem Sets 3

Sept. 24, 2026

Problem 1. Let I be an open interval in $\mathbb{R}$, $c \in I$, and $f : I \rightarrow \mathbb{R}$ be a function. Show that f is continuous at c if and only if $\lim_{h \rightarrow 0} f(c+h) = f(c)$.

Problem 2. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function satisfying $f(a+b) = f(a)f(b)$ for all $a, b \in \mathbb{R}$.

1. Show that $f(x) \geq 0$ for all $x \in \mathbb{R}$.
2. Show that if f is continuous at 0, then f is continuous on $\mathbb{R}$ (that is, f is continuous at every point of $\mathbb{R}$).

Problem 3. Let I be an interval in $\mathbb{R}$ and $f, g : I \rightarrow \mathbb{R}$ be continuous functions. Show that if $f(x) = g(x)$ for all $x \in \mathbb{Q} \cap I$, then $f(x) = g(x)$ for all $x \in I$.

Problem 4. Let I be an interval, $c \in I$, and $f : I \rightarrow \mathbb{R}$ be a continuous function. Show that if $f(c) \neq 0$, there exists $\delta > 0$ such that $f(x)f(c) > 0$ whenever $|x - c| < \delta$ and $x \in I$.

Problem 5. Construct a function $f : \mathbb{R} \rightarrow \mathbb{R}$ so that f is continuous at all integers but nowhere else.

Problem 6. Show that the equation $x^3 - 15x + 1 = 0$ has three solutions in the interval $[-4, 4]$.

Problem 7. Suppose that a and b are positive constants. Show that the equation

$$\frac{a}{x^3 + 2x^2 - 1} + \frac{b}{x^3 + x - 2} = 0$$

has at least one solution in the interval $(-1, 1)$.

Problem 8. True or False: Determine whether the following statements are true or false. If it is true, prove it. Otherwise, give a counter-example.

1. If f and g are functions such that $\lim_{x \rightarrow c^+} g(x) = K$, $\lim_{y \rightarrow K^+} f(y) = f(K)$, then

$$\lim_{x \rightarrow c^+} (f \circ g)(x) = f(K).$$

How about if $x \rightarrow c^+$ and $y \rightarrow K^+$ are replaced by $x \rightarrow c^-$ and $y \rightarrow K^-$, respectively?

2. If $|f|$ is continuous at c , so is f .
3. Let I be an interval and $f : I \rightarrow \mathbb{R}$ be a continuous function. If $f(x) \neq 0$ for all $x \in I$, then f never change signs; that is, either $f(x) > 0$ for all $x \in I$ or $f(x) < 0$ for all $x \in I$.
4. If $\lim_{x \rightarrow c} f(x) = \infty$ and $\lim_{x \rightarrow c} [f(x) - g(x)] = 0$, then $\lim_{x \rightarrow c} g(x) = \infty$.